

# GROWTH AND ISOPERIMETRIC PROFILE OF PLANAR GRAPHS

ITAI BENJAMINI AND PANOS PAPASOGLU

**ABSTRACT.** Let  $\Gamma$  be a planar graph such that the volume function of  $\Gamma$  satisfies  $V(2n) \leq CV(n)$  for some constant  $C > 0$ . Then for every vertex  $v$  of  $\Gamma$  and  $n \in \mathbb{N}$ , there is a domain  $\Omega$  such that  $B(v, n) \subset \Omega$ ,  $\partial\Omega \subset B(v, 6n)$  and  $|\partial\Omega| \lesssim n$ .

## 1. INTRODUCTION

Let  $\Gamma$  be a locally finite graph. If  $a$  is a vertex of  $\Gamma$  we denote by  $B(a, n)$  the ball of radius  $n$  centered at  $a$ . If  $B$  is a graph, we denote by  $|B|$  the number of vertices of  $B$ .

Define the *growth function* of  $\Gamma$  at  $a$  by

$$V(a, n) = |B(a, n)|$$

A graph  $\Gamma$  is *doubling* if there is a constant  $C > 0$  such that for all  $a, b \in \text{Vert}(\Gamma)$  and  $n \in \mathbb{N}$ ,  $V(a, 2n) \leq CV(b, n)$ . We say then that  $C$  is a *doubling constant* for  $\Gamma$ .

If  $\Gamma$  is a doubling graph then the degree of vertices is uniformly bounded.

Note that for any  $d \geq 1$ , there are planar graphs with the doubling property for which, for any  $a$  and  $n$ ,  $V(a, n)$  is of order  $n^d$ , see e.g. the last section of [4].

Say that a graph  $\Gamma$  corresponds to a *tessellation* of  $\mathbb{R}^2$  if there is a  $k \in \mathbb{N}$  such that all components of  $\mathbb{R}^2 - \Gamma$  are bounded regions with at most  $k$  sides.

**Definition .** Let  $(X, d)$  be a metric space. An  $\epsilon$ -*net*  $N$  of  $X$  is a set such that  $d(v_1, v_2) > \epsilon$  for all  $v_1, v_2 \in N$  and  $N$  is maximal set with this property.

We remark that if  $N$  is an  $\epsilon$ -net of  $X$  then  $X$  is contained in the  $\epsilon$ -neighborhood of  $N$ .

**Definition .** Let  $\Omega$  be a subgraph of a graph  $\Gamma$ . Then we denote by  $\partial\Omega$  the set of all vertices of  $\Gamma - \Omega$  which have a neighbor in  $\Omega$ .

---

1991 *Mathematics Subject Classification.* 05C10, 53C20, 53C23.

Our main result is the following:

**Theorem 2.1.** *Let  $\Gamma$  be a doubling planar graph. Then there is a constant  $\alpha$  so that for every vertex  $v$  of  $\Gamma$  there is a finite domain  $\Omega$  such that  $B(v, n) \subset \Omega$ ,  $\partial\Omega \subset B(v, 6n)$  and  $|\partial\Omega| \leq \alpha n$ .*

Krikun [7] has shown a similar theorem for the uniform infinite planar triangulation (UIPT) introduced in [2]. The volume doubling property does not hold for the UIPT, still an asymptotic version should hold: for any vertex  $v$ , for large enough  $n$ ,  $B(v, 2n)$  contains order 1 disjoint balls of radius  $n/2$  a.s. and the proof below will adapt to give Krikun's result. The asymptotic volume growth of balls in the UIPT is order  $n^4$ , up to polylog's, see [1], thus a weaker result with a polylog correction follows from our result. For the uniform infinite planar quadrangulation a sharp volume growth estimate is known which implies the asymptotic volume doubling [8, 5]. For a detailed study of the geometry of the uniform infinite planar quadrangulation and a matching lower bound on the small cuts see [5].

Our result shows that the volume and the isoperimetric profile function are related for planar graphs. We define here the isoperimetric profile function of a graph in a similar way as for Riemannian manifolds:

**Definition .** Let  $\Gamma$  be a locally finite graph and let  $V(n)$  be the volume function of  $\Gamma$ . Then the *isoperimetric profile* function of  $\Gamma$ ,  $I_\Gamma : \mathbb{N} \rightarrow \mathbb{N}$  is defined by:

$$I_\Gamma(n) = \inf_{\Omega} \{|\partial\Omega| : \Omega \subset \Gamma, |\Omega| \leq n\}$$

where  $\Omega$  ranges over all subgraphs of  $\Gamma$ .

From theorem 2.1 we obtain the following:

**Corollary 1.1.** *Let  $\Gamma$  be a doubling planar graph with volume function  $V(n)$  and isoperimetric profile function  $I_\Gamma(n)$ .*

*Let  $\varphi(n) = \inf\{k : V(k) \geq n\}$ . Then there is a constant  $\alpha$  such that*

$$I_\Gamma(n) \leq \alpha \varphi(n) \text{ for all } n \in \mathbb{N}$$

The same result holds for Riemannian metrics on the plane or on the plane with holes (again assuming the doubling property holds for the metric) and our proof extends to this setting as well (see [9] for more information on isoperimetric profiles of planes and planes with holes).

**1.1. Idea of the proof.** Here is a sketch of the proof of theorem 2.1. Let  $v$  be any vertex of  $\Gamma$ . Consider the balls  $B(v, n), B(v, 3n)$ . Let  $N$  be an  $n$ -net of  $\partial B(v, 3n)$ . For each vertex  $w$  of  $N$  consider  $B(w, n/2)$ . Note that all such balls are disjoint since  $N$  is an  $n$ -net. Also all these balls are contained in  $B(v, 4n)$ . So, by the doubling property, we can have only boundedly many such balls, that is  $|N| \leq \beta$ , where  $\beta$  does not depend on  $n$ . Consider now the balls  $B(w, 2n)$  for all  $w \in N$ .  $\partial B(v, 3n)$  is contained in the union of these balls. Construct a closed curve that ‘blocks’  $v$  from infinity as follows: if  $w_1, w_2 \in N$  are such that  $d(w_1, w_2) \leq 2n$  then we join them by a geodesic. So replace  $\partial B(v, 3n)$  by the ‘polygonal line’ that we define using vertices in  $N$ . This ‘polygonal line’ blocks  $v$  from infinity and has length at most  $2n\beta$ . In the next section we make precise this idea. There are some technical issues to take care of, for example  $\partial B(v, 3n)$  might not be connected (and could even have ‘large gaps’) and the geodesic segments have to be chosen carefully. In particular the constants obtained will be slightly different from the ones in this sketch.

## 2. GROWTH AND PROFILE

**Definition .** Let  $\Gamma$  be an infinite planar locally finite graph. We say that an embedding of  $\Gamma$  in the plane is *tame* if for any bounded subset  $A$  of the plane,  $\Gamma - A$  has at least one connected component of infinite diameter. It is easy to see that any infinite planar locally finite graph admits a tame embedding to  $\mathbb{R}^2$ .

We see now  $\Gamma$  as embedded in the plane by a tame embedding. Let  $B = B(v, r)$  be a ball of  $\Gamma$  and let  $U$  be the unbounded connected component of  $\mathbb{R}^2 - B$ . Define the *contour* of  $B(v, r)$  to be the graph  $B \cap \bar{U}$ . Note that the contour of  $B(v, r)$  is connected.

Recall some basic facts about winding numbers (see e.g. [6], ch. 3 for a definition and basic properties of winding numbers).

Let  $\gamma$  be a closed curve on the plane and  $v$  be a point that does not lie on  $\gamma$ . If the winding number  $W(\gamma, v)$  is non zero then  $v$  lies in a bounded component of  $\mathbb{R}^2 - \gamma$ . If  $\gamma = \gamma_1 \cup \gamma_2$  where  $\gamma_1, \gamma_2$  are paths with the same endpoints  $a, b$  and if  $\delta$  is another path with endpoints  $a, b$  then

$$W(\gamma, v) = W(\gamma_1 \cup \delta, v) + W(\gamma_2 \cup \delta, v)$$

If  $\gamma$  is a closed curve then  $W(\gamma, v) \in \mathbb{Z}$ . Here it will be convenient to consider the winding number modulo 2, so in what follows by winding number of  $\gamma$  around  $v$  we mean  $W(\gamma, v) \bmod 2$ .

**Theorem 2.1.** *Let  $\Gamma$  be a doubling planar graph. Then there is a constant  $\alpha$  so that for every vertex  $v$  of  $\Gamma$  there is a finite domain  $\Omega$  such that  $B(v, n) \subset \Omega$ ,  $\partial\Omega \subset B(v, 6n)$  and  $|\partial\Omega| \leq \alpha n$ .*

*Proof.* Consider a tame embedding of  $\Gamma$  in the plane. Let  $F$  be the contour of  $B(v, 4n)$  with respect to this embedding. Clearly  $F$  is a connected subset of the plane. ‘Parametrise’  $F$  by a map  $f : S \rightarrow F$  where  $S$  is a graph homeomorphic to the circle,  $f$  sends edges to edges and every vertex  $w$  such that  $d(w, v) = 4n$ , has a unique pre-image  $f^{-1}(w)$ . Note that  $F$  is not necessarily a simple closed curve, for example consider the case that  $\Gamma$  is a tree. Denote by  $B'$  the set of all vertices  $w$  in  $\Gamma$  such that  $d(w, v) = 4n$ .

It is possible that  $v \in F = \text{im } f$ . As this creates some technical problems we modify  $f$  slightly to avoid this, so that the curve  $f(S)$  goes around  $v$ . For example we may do this as follows: Cyclically order the edges adjacent to  $v$ , as  $e_1 = [v, v_1], \dots, e_k = [v, v_k]$  and introduce new edges  $e'_1, \dots, e'_k$  so that the edge  $e'_i$  joins  $v_i, v_{i+1}$  ( $i \in \mathbb{Z}_k$ ). One may assume that the  $e'_i$ s do not intersect the interior of any edge of  $\Gamma$  and that the only edges contained in the interior of  $e'_1 \cup \dots \cup e'_k$  are  $e_1, \dots, e_k$ .

If for two successive edges  $g_1, g_2$  of  $S$ , we have  $f(g_1) = e_i, f(g_2) = e_{i+1}$  modify  $f$  so that  $f(g_1 \cup g_2) = e'_i$ .

After this modification the winding number of  $f(S)$  around  $v$  is defined and it is equal to 1. If  $S_1$  is a subarc of  $S$  define the *weight* of  $S_1$  to be  $w(S_1) = |f(S_1) \cap B'|$ .

Note that if  $a, b \in B'$  then there are two subarcs  $S_1, S_2$  of  $S$  such that  $S = S_1 \cup S_2$  and  $f(S_1), f(S_2)$  are paths with endpoints  $a, b$ . If we further assume that  $d(a, b) < 4n$  and  $\gamma$  is a geodesic in  $\Gamma$  joining  $a, b$  then  $v \notin \gamma$ , so one of the closed curves  $f(S_1) \cup \gamma$  and  $f(S_2) \cup \gamma$  has winding number 1 around  $v$ , while the other has winding number 0 around  $v$ .

If for any two vertices  $a, b$  of  $B'$ ,  $d(a, b) > 2n + 1$  then the balls  $B(a, n)$ , where  $a \in B'$  are disjoint. So by the doubling property  $|B'| \leq C^3$ , where  $C$  is the doubling constant of  $\Gamma$ . Therefore in this case one can take  $\Omega = B(v, 4n)$ .

Otherwise pick  $a, b$  in  $B'$  with the following two properties:

1)  $d(a, b) \leq 2n + 1$ .

2) There is a geodesic  $\gamma$  in  $\Gamma$  joining  $a, b$  so that for the subarc  $S_1 \subset S$  for which the closed curve  $f(S_1) \cup \gamma$  has winding number 1 around  $v$ ,  $w(S_1)$  is minimum possible.

If  $f(S_1) \cap B' = \{a, b\}$  we may take  $\Omega$  to be the component of  $\Gamma - \gamma$  containing  $v$  and the theorem is proved.

Otherwise, observe that  $f(S_1) \cup \gamma$  separates  $v$  from infinity. We now explain how to replace  $f(S_1)$  by a curve of length linear in  $n$ .

Consider an  $n$ -net  $N$  of the set  $B' \cap f(S_1)$ . Observe that if  $x, y \in N$  the balls  $B(x, n/2), B(y, n/2)$  are disjoint and are contained in  $B(v, 9n/2)$ . It follows that  $|N| < C^4$ .

Define a graph  $\Delta$  with set of vertices equal to  $N \cup \{a, b\}$ . Join two vertices  $x, y$  by an edge if  $d(x, y) \leq 2n + 1$ .

We consider first the case that  $\Delta$  is connected. Then there is a simple path in  $\Delta$  joining  $a, b$ . This simple path corresponds to a path  $\delta$  in  $\Gamma$ . We obtain  $\delta$  by replacing edges of the path in  $\Delta$  by geodesic paths of length  $\leq 2n + 1$  in  $\Gamma$ . It follows that

$$\text{length}(\delta) \leq (2n + 1)C^4.$$

Consider now the closed path

$$p = \gamma \cup \delta$$

We claim that the winding number of  $p$  around  $v$  is 1. Indeed  $\delta = \delta_1 \cup \dots \cup \delta_k$  where  $\delta_i = [a_i, a_{i+1}]$  are geodesic paths of  $\Gamma$  corresponding to edges of  $\Delta$  as explained above and  $a = a_1, b = a_{k+1}$ . Clearly there are  $a'_1, \dots, a'_{k+1} \in S_1$  such that  $f(a'_i) = a_i$  and

$$\bigcup_{i=1}^k (a'_i, a'_{i+1}) = S_1.$$

The curve

$$f(a'_i, a'_{i+1}) \cup \delta_i$$

is closed and has winding number 0 around  $v$ . This follows from the definition of  $\gamma$  and the fact that the weight of  $(a'_i, a'_{i+1})$  is smaller than the weight of  $S_1$ . By the additivity property of winding numbers we see that by replacing successively  $f(a'_i, a'_{i+1})$  by  $\delta_i$  in  $f(S_1) \cup \gamma$  the winding number around  $v$  remains the same. So the winding number of  $p$  around  $v$  is 1 and the claim is proved.

Let  $\Omega$  be the component of  $\Gamma - p$  containing  $v$ .  $\Omega$  clearly has all the properties required by the theorem.

Let's return now to the proof of the general case, so we don't assume anymore that  $\Delta$  is connected.

Let  $\Delta_1$  be the connected component of  $\Delta$  containing  $a$ . Let  $V_1$  be the vertex set of  $\Delta_1$ .

Let's denote by  $a', b'$  the endpoints of  $S_1$ , so  $f(a') = a, f(b') = b$ .

Let  $a'_2 \in S_1$  be such that

$$d(f(a'_2), V_1) \leq n$$

and  $f[a', a'_2]$  contains all vertices of  $f(S_1)$  that are at distance  $\leq n$  from  $V_1$ . Join  $f(a'_2)$  to a vertex of  $\Delta_1$  by a geodesic  $\gamma_1$  of length  $\leq n$ . If  $d_1 = |V_1|$  then, as before, we see that there is a path  $\delta_1$  joining  $a$  to  $f(a'_2)$  such that

$$\text{length}(\delta_1) \leq d_1(2n + 1)$$

and the winding number of  $\gamma \cup \delta_1 \cup f[a'_2, b']$  around  $v$  is 1.

Let  $b'_2$  be the first vertex of  $S_1$  following  $a'_2$  such that  $d(f(b'_2), v) = 4n$ . Consider an  $n$ -net  $N_1$  of the set  $B' \cap f(a'_2, b')$ . Clearly  $N_1 \cup V_1$  is an  $n$ -net of the set  $B' \cap f(S_1)$  so  $|N_1 \cup V_1| \leq C^4$ . As before define a graph with vertex set  $\{f(b'_2)\} \cup N_1$  and consider the connected component, say  $\Delta_2$ , of this graph which contains  $f(b'_2)$ . If  $V_2$  is the vertex set of  $\Delta_2$ , let  $a'_3 \in S_1$  be such that

$$d(f(a'_3), V_2) \leq n$$

and  $f[b'_2, a'_3]$  contains all vertices of  $f(S_1)$  that are at distance  $\leq n$  from  $V_2$ . Join  $f(a'_3)$  to a vertex of  $\Delta_1$  by a geodesic  $\gamma_2$  of length  $\leq n$ . If  $d_2 = |V_2|$  then, as before, we see that there is a path  $\delta_2$  joining  $f(b'_2)$  to  $f(a'_3)$  such that

$$\text{length}(\delta_2) \leq d_2(2n + 1)$$

and the winding number of

$$\gamma \cup \delta_1 \cup f[a'_2, b'_2] \cup \delta_2 \cup f[a'_3, b']$$

around  $v$  is 1.

We continue in the same way. Consider an  $n$ -net  $N_2$  of  $f[a'_3, b'] \cap B'$ , define similarly a graph  $\Delta_3$  and a path  $\delta_3$ . Clearly this procedure terminates and eventually we produce paths  $\delta_1, \dots, \delta_k$  such that:

1) The winding number of

$$\gamma \cup \delta_1 \cup f[a'_2, b'_2] \cup \dots \cup f[a'_k, b'_k] \cup \delta_k$$

around  $v$  is 1.

2)  $f(a'_i, b'_i) \cap B' = \emptyset$  for all  $i = 2, \dots, k$ .

3)  $\sum_{i=1}^k \text{length}(\delta_i) \leq C^4(2n + 1)$ .

4) The paths  $\gamma, \delta_1, \delta_2, \dots, \delta_k$  are contained in  $B(v, 6n) - B(v, n)$ .

Let  $\Omega$  be the connected component of  $\Gamma - (\gamma \cup \delta_1 \cup \dots \cup \delta_k)$  containing  $v$ . By the definition of  $\Omega$ ,  $B(v, n) \subset \Omega$ . By property 3 above  $|\partial\Omega| \leq (C^4 + 1)(2n + 1)$ .

□

### Remarks:

1. If  $\Gamma$  corresponds to a tessellation then, the proof of the theorem shows that there is a simple closed curve of length  $\sim n$  in  $B(v, 6n)$  that separates  $B(v, n)$  from infinity. In the random context this has been

shown by Krikun ([7], [8]) for triangulations and quadrangulations. Our proof applies regardless of the ‘shape’ of the regions of the tessellation and with uniform bounds.

2. Given any  $\epsilon > 0$  one may easily adapt the proof of the theorem to produce regions  $\Omega$  with the same properties such that

$$\partial\Omega \subset B(v, (1 + \epsilon)n)$$

for  $n$  sufficiently large. Of course in this case one obtains  $|\partial\Omega| \leq C(\epsilon)n$  and  $C(\epsilon) \rightarrow \infty$  as  $\epsilon \rightarrow 0$ .

### 3. FURTHER COMMENTS

A graph  $G$  admits uniform volume growth  $f(n)$ , if there are  $0 < c < C < \infty$ , so that for all  $n$ , any ball of radius  $n$  in  $G$  satisfies,

$$cf(n) < |B(v, n)| < Cf(n).$$

For planar graph admitting arbitrarily large uniform polynomial growth see e.g. [4]. It is conjectured that planar graphs of uniform polynomial growth are recurrent for the simple random walk. It is also conjectured [2] that the UIPT is recurrent, see also [4]. By the Nash-Williams sufficient condition for recurrence, it is enough to find infinitely many disjoint cutsets  $\{C_i\}$  separating the root from infinity. so that  $\sum |C_i|^{-1} = \infty$ . The theorem above is a step in that direction, still we don’t know if planar graphs of uniform polynomial growth admits such cutsets? Maybe not.

Assume  $G$  is a planar graph of uniform polynomial growth  $n^d, d > 2$ , by the theorem above  $G$  admits bottlenecks. This suggests the *conjecture* that simple random walk on  $G$  will be subdiffusive, as it will spend a lot of time in domains with small boundary before exiting. That is, the expected distance to the root by time  $t$  is bounded by  $t^\alpha$  for  $\alpha < 1/2$ . Does  $\alpha = d^{-1}$ ? Subdiffusivity was recently established for the UIPT by Nicolas Curien and the first author.

What about a high dimensional generalization? A  $d$ -sphere packing is a collection of  $d$ -dimensional balls with disjoint interiors. Associated to the packing an unoriented graph  $G = (V, E)$  called the *d-tangency graph*, where vertices corresponds to the  $d$ -balls and edges are between any two tangent balls, see [3]. Is it the case that for any  $d$  a  $d$ -tangency graph with the doubling property admits cutsets outside a ball of radius  $n$  of size  $n^{d-1}$ ?

Let  $G$  be a planar triangulation which is doubling and further assume all balls has growth  $r^d, d > 2$  up to a multiplicative constant. Is there such  $G$  for which all complements of balls are connected, for all

balls? Or as in the UIPT, the complements of some balls admit several connected components, some of size proportional to the ball?

## REFERENCES

- [1] O. Angel, Growth and percolation on the uniform infinite planar triangulation, *Geometric And Functional Analysis* **13** 935–974 (2003).
- [2] O. Angel and O. Schramm, Uniform infinite planar triangulations, *Comm. Math. Phys.* **241**, 191–213 (2003).
- [3] I. Benjamini and N. Curien, Local limit of Packable graphs, <http://arxiv.org/abs/0907.2609>
- [4] I. Benjamini and O. Schramm, Recurrence of Distributional Limits of Finite Planar Graphs, *Electron. J. Probab.* **6** 13 pp. (2001).
- [5] N. Curien, L. Menard and G. Miermont, The uniform infinite planar quadrangulation seen from infinity. In preparation.
- [6] W. Fulton, Algebraic Topology, Springer 1995.
- [7] M. Krikun, Uniform infinite planar triangulation and related time-reversed critical branching process, arXiv:math/0311127
- [8] M. Krikun, Local structure of random quadrangulations, arXiv:math/0512304
- [9] P. Papasoglu, *Cheeger constants of surfaces and isoperimetric inequalities*, Trans. Amer. Math. Soc. 361 (2009), 5139–5162.

*E-mail address:* itai.benjamini@weizmann.ac.il

*E-mail address:* papazoglou@maths.ox.ac.uk

DEPT. OF MATHEMATICS, WEIZMANN INSTITUTE, REHOVOT, 76100, ISRAEL

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, 24-29 ST GILES', OXFORD, OX1 3LB, U.K.